

Case Report

A Heavy Infection Macro-cysts of *Sarcocystis* Spp. Case in a Najdi Sheep at Buraidah Slaughter House, Qassim, Saudi Arabia

Mohammad Ali Al-wabel, Shawkat Mohamed Fat'hi*

College of Agriculture and Veterinary Medicine, Qassim University, Buriadah, Saudi Arabia

Email address:

shawkatfathi@gmail.com (S. M. Fat'hi), sabdellatie@qu.edu.sa (S. M. Fat'hi)

*Corresponding author

To cite this article:Mohammad Ali Al-wabel, Shawkat Mohamed Fat'hi. A Heavy Infection Macro-cysts of *Sarcocystis* Spp. Case in a Najdi Sheep at Buraidah Slaughter House, Qassim, Saudi Arabia. *Animal and Veterinary Sciences*. Vol. 8, No. 5, 2020, pp. 96-98. doi: 10.11648/j.avs.20200805.11**Received:** August 1, 2020; **Accepted:** August 12, 2020; **Published:** August 31, 2020

Abstract: Background: Protozoal diseases caused by *Sarcocystis* species can cause serious damage in sheep and goats flocks, inducing decreased growth conversion rates and partial or complete condemnation of carcasses at the slaughter house. *Sarcocystis* spp. affecting sheep and goats are host specific for their intermediate hosts and family specific for their final hosts. They derive their name from the intramuscular cyst stage present in the intermediate host. Objective: Macro and micro identification of macro-cysts of *Sarcocystis* spp. affecting a Najdi sheep carcass. Method: Detection of macrosarcocystis during routine post mortem examination and identification the detected cyst by histopathological investigation. Result: The macroscopic sarcocystis were found cleared and scattered in almost all skeletal muscles including shoulder, chest and diaphragm as well as thigh. Both morphological and histopathological characteristics indicated that the detected cysts may be identified as *Sarcocystis gigantea*. Conclusion: It is a case study of macro-cysts of *Sarcocystis* spp. affecting local breed of Najdi sheep. It is considered a seldom detected case all over many years during meat inspection at Buraidah slaughter house. The possible control measures are those of simple hygiene. Farm dogs should not be housed in, or allowed access to, fodder stores nor should they be allowed to defecate in pens where livestock are housed. It is also important that they are not fed uncooked meat.

Keywords: Najdi Sheep, Macroscopic *sarcocystis*, Histopathology

1. Introduction

Sarcocystis spp. are cyst-forming coccidia that infect several livestock species causing sarcocystosis which responsible for the formation of cysts often observed in the skeletal and cardiac muscles. It has a high prevalence in sheep, resulting in serious injury in different organ systems, abortion, and death. There are two common *Sarcocystis* species affecting sheep, the first is *Sarcocystis tenella* (*S. ovis*) which is highly pathogenic to lambs causing fever, anorexia, weight loss, loss of wool, abortion, premature birth nervous signs and death. The Cysts are microscopic and located in the muscles of oesophagus, diaphragm, heart, tongue and central nervous system. The second species is *Sarcocystis gigantea* (*S. ovifelis*) which is moderately pathogenic to sheep with

macroscopic cysts found as small white cysts resemble grains of rice and commonly affect the oesophagus, tongue, diaphragm and skeletal muscle and their presence in the muscles and viscera of carcasses at the slaughterhouse causes economic losses due to the partial or total loss of carcasses. The sarcocystosis of sheep requires two hosts to complete its life cycle, where the cat acts as its definitive host in case *S. ovifelis*. Cats become infected when they feed on infected sheep carcasses. Ingested *sarcocystis* cysts develop in the cat to produce tiny spores which passed in cat faeces contaminating pasture and/or stock feed. Sheep is considered intermediate host which ingest *sarcocystis* spores when feeding on contaminated pasture or hay/grain. Spores move from the gut to small blood vessels where they multiply and move into muscle tissue and develop into cysts [1-5].

Identification of *sarcocystis* spp. in sheep is done by

morphological, ultrastructural, and molecular tests [2]. Morphologically, *sarcocystis* cysts appear as visible elongated cylindrical bodies in striated muscle and sometimes in unstriated muscle [6]. They occur as spindle to oval in shape and two stages were recognized, the peripheral metrocytes and centrally located banana-shaped bradyzoites [7]. Light microscopy [7] and Electron Microscopy (EM) [8] were applied for ultrastructural characteristics of *sarcocystis* spp. In Saudi Arabia, macroscopic cysts were usually detected on the oesophagus, where EM showed an ultrastructural similarities in the cyst walls and their parasite contents (metrocytes and merozoites) for all macroscopic cysts examined from sheep and goats. These were also enclosed by a secondary cyst wall. The primary cyst wall had many cauliflower-like protrusions supported by many fibellar structures, while the ground substance underneath them contained many dark, dense granules and usually extended into the cyst cavity dividing it into many compartments containing the parasites. The cyst parasites were the peripheral globular to spherical metrocytes, which were divided by endodyogeny and the central banana-shaped merozoites. Both metrocytes and merozoites had all characteristics of the Apicomplexa [8]. Furthermore, identification of *sarcocystis* spp. in sheep by PCR products of the 18S rRNA gene were successfully sequenced [9-11].

2. Materials and Methods

2.1. Meat Inspection

The routine post mortem examination of a Najdi sheep carcass revealed multiple macroscopic *sarcocystis* cysts. The tissue samples were collected from different affected muscles by *sarcocystis* macrocysts for histopathological examination.

2.2. Histopathological Investigation

Tissue specimens were preserved in 10% neutral buffered

formalin immediately after removal from the carcass for fixation. For routine paraffin wax histopathology, a portion of 1cm³ lesion from each of the affected muscle was incised, fixed in 10% formal saline, processed and stained with H&E as described by [12-14].

3. Results and Discussion

Sarcocystosis is a parasitic disease caused by varying *Sarcocystis* species infecting humans and animals. *Sarcocystis* spp. are cyst-forming coccidia that infect numerous animals species, including several livestock species. It is commonly found in small ruminants causing pathogenic effects [11].

Multiple macroscopic sarcocystis were observed and distributed randomly throughout nearly all the muscles of a famous domestic Najdi sheep at Qassim Region, Saudi Arabia. They characterized by white, round, oval cysts or banana-shaped bradyzoites as demonstrated in figure 1 and described by [7].

The histomorphological features of both longitudinal and cross muscular sections of sarcocystosis were illustrated in "Figure 2". The longitudinal sections showed spindle shaped encysted protozoan (arrow) with well-defined wall containing several separated bradyzoites (arrow head), muscular degeneration, fat depositing (asterisk) as a result of necrosis and atrophy and mild inflammatory cells infiltrates "Figures 2, A & B". While, the cross muscular sections revealed sarcocysts with a well-defined wall (arrow head) containing several separated bradyzoites with centric pale halo (asterisk) and eosinophilic degenerative muscle fibers (arrow) "Figures 2, C & D".

The morphology of macroscopic sarcocystis found in the muscles of such case mostly resembled that of the *S. gigantea* cysts, where the diagnosis was based on the morphological as well as histopathological findings.




Figure 1. Chest and diaphragm muscular macroscopic sarcocystosis of a Najdi sheep carcass.

Sarcocystis muscular cysts are varying from microscopic to macroscopic size. Sheep can be infected by four species of Sarcocystis including *S. tenella*, *S. arieticanis*, *S. gigantea* and *S. medusiformis*. *S. tenella* and *S. arieticanis* are known as pathogenic parasites, which form muscular micro-cysts transferred by dogs, while *S. gigantea* and *S. medusiformis* are non-pathogenic and form macro-cysts transferred by cats [2, 10].

Further identification of either microscopic or macroscopic sarcocystis cysts affection in sheep can be performed by molecular assays [15].




Figure 2. Histomorphology of muscular sarcocystosis (H&E, 10X and 40X).

References

- [1] Martínez-Navalón, B., B. Anastasio-Giner, M. Cano-Fructuoso, P. Sanchez-Martínez, A. Llopis-Morant, B. Perez-Castarlenas, E. Goyena, E. Berriatua Fernández de Larrea, 2012. Short communication. Sarcocystis infection: a major cause of carcass condemnation in adult sheep in Spain. Spanish Journal of Agricultural Research, v. 10, n. 2, p. 388-392, 2012.
- [2] Bittencourt, Marta Vasconcelos; Iris Daniela S. Meneses; Müller Ribeiro-Andrade; Rogério Fernando de Jesus; Flávio Ribeiro de Araújo; Luís F. Pita Gondim, 2016. Sarcocystis spp. in sheep and goats: frequency of infection and species identification by morphological, ultrastructural, and molecular tests in Bahia, Brazil. Parasitology Research, Vol. 115, Issue 4, pp 1683–1689.
- [3] Dubey JP, Calero Bernal R, Rosenthal BM, Speer CA. Fayer. Sarcocystosis of animals and humans. 2nd Edn. Boca Raton: CRC Press; Taylor & Francis Group; 2016.
- [4] Asmaa, Dahmani, Harhoura Khaledb, Aissi Miriemb, Zenia Safiab, Saadi Ahmedc, Kadour Rachid, 2017. Study of ovine sarcosporidiosis in slaughterhouses of El Harrach in north of Algeria. Veterinaria, Vol. 66, No. 3, 2017.
- [5] Dong, Hui, Ruijing Su, Yinghua Wang, Zongxi Tong, Longxian Zhang, Yurong Yang and Junjie Hu, 2018. Sarcocystis species in wild and domestic sheep (*Ovis ammon* and *Ovis aries*) from China. BMC Veterinary Research, 14: 377, 1-7.
- [6] Dehaghi, MM; Fallahi, M; Sami, M; Radfar, MH, 2013. Survey of sarcocystis infection in slaughtered sheep in Kerman Abattoir, Kerman, Iran. Comparative Clinical Pathology, May 2013, Vol. 22, Issue 3, pp 343–346.
- [7] Latif B, Kannan Kutty M, Muslim A, Hussaini J, Omar E, Heo CC, Rossle NF, Abdullah S, Kamarudin MA, Zulkarnain MA., 2015. Light microscopy and molecular identification of Sarcocystis spp. in meat producing animals in Selangor, Malaysia. Trop Biomed. 2015 Sep; 32 (3): 444-52.
- [8] Al-Hoot AS, Al-Qureishy SA, Al-Rashid K, Bashtar AR., 2005. Microscopic study on Sarcocystis moulei from sheep and goats in Saudi Arabia. J Egypt Soc Parasitol. 2005 Apr; 35 (1): 295-312.
- [9] da Silva RC, Su C, Langoni H., 2009. First identification of Sarcocystis tenella (Railliet, 1886) Moulé, 1886 (Protozoa: Apicomplexa) by PCR in naturally infected sheep from Brazil. Wet Parasitol 2009 Nov 12; 165 (3-4): 332-6.
- [10] Kalantari, N., Mohaddeseh Khaksar, Salman Ghaffari, and Seyed Mehdi Hamidekish, 2016. Molecular Analysis of Sarcocystis Spp. Isolated from Sheep (*Ovis aries*) in Babol Area, Mazandaran Province, Northern Iran. Iran J Parasitol. 2016 Jan-Mar; 11 (1): 73–80.
- [11] Amairia S, Amdouni Y, Rouatbi M, Rjeibi MR, Awadi S, Gharbi M., 2018. First detection and molecular identification of Sarcocystis spp. in small ruminants in North-West Tunisia. Transbound Emerg Dis. 2018 Apr; 65 (2): 441-446.
- [12] Crocker, J. (2005). Histopathology section in: The Science of laboratory, 2nd ed. Pp. 566.
- [13] Hewitson, T. D. & Dabry, I. A. (2010). Histology Protocols, Human press, Pp: 225.
- [14] Kidsadagon P. (2012). Field necropsy techniques in mammal and poultry. Faculty of Veterinary medicine, Chiang Mai University manual, pp: 41-50.
- [15] Bahari, P; Mitra Salehi, Mohsen Seyedabadi, and Ahmad Mohammadi, 2014. Molecular Identification of Macroscopic And Microscopic Cysts of Sarcocystis in Sheep in North Khorasan Province, Iran. Int J Mol Cell Med. 2014; 3 (1): 51–56.